

**BRUMOIDES SEPTENTRIONIS DAVISI (LENG)
(COLEOPTERA: COCCINELLIDAE): DISTRIBUTION,
HOST-PLANT ASSOCIATIONS, AND HABITATS OF A
SELDOM-COLLECTED LADY BEETLE**

A. G. WHEELER, JR.

Department of Entomology, Clemson University, Clemson, SC 29634, U.S.A. (e-mail: awhlr@clemson.edu)

Abstract.—The chilocorine coccinellid *Brumoides septentrionis davisii* (Leng) is an infrequently collected lady beetle whose broad distribution in eastern North America has been mapped; few specific localities, however, have been published. Despite a known association with pines, the *Pinus* species on which it is found have not been recorded. Historical locality data are provided based on specimens in five museum collections, and new records from recent field work are presented. This coccinellid's plant and habitat associations are discussed. Its populations apparently have declined in the Northeast in recent years. Certain scale insects probably serve as prey.

Key Words: *Brumoides septentrionis davisii*, coccinellids, insect distribution, pine barrens

The most frequently encountered of all predacious beetles (Clausen 1940), coccinellids are associated with biological control more than any other group of predators (Obrycki and Kring 1998). Coccidophagous lady beetles have proved more effective as classical biological control agents than those that are aphidophages (Dixon 2000). The establishment, either from bio-control introductions or accidentally from commerce, of several Old World coccinellids that are generalist aphid predators has led to more stringent regulations for their importation and has prompted interest in native lady beetles whose densities have declined as those of adventive coccinellids have increased (e.g., Schaefer et al. 1987, Wheeler and Hoebeke 1995, Colunga-Garcia and Gage 1998, Cottrell and Yeargan 1998, Obrycki et al. 1998, Evans 2000).

I provide distributional records of the Nearctic *Brumoides septentrionis davisii*

(Leng) and notes on its plant associations and habitats to call attention to this infrequently collected lady beetle. I also suggest possible prey associations and a recent decline in its numbers. The availability of this background information might prove useful if studies on the bionomics of this coccinellid are initiated.

Brumoides septentrionis davisii (Leng)

The chilocorine *Brumoides septentrionis davisii* (*sensu* Gordon 1985) is the eastern North American subspecies of *B. septentrionis* (Weise), whose other subspecies are found farther west in Canada (McNamara 1991) and the United States (Gordon 1985). Leng (1908) described the eastern subspecies as *Exochomus (Brumoides) septentrionis* var. *davisii* from Alabama, District of Columbia, "Lake Superior," Massachusetts, Michigan, New Jersey, Pennsylvania, and Virginia; except for Lakehurst, N.J., specific localities were not mentioned. The sub-

genera of *Exochomus* that Leng (1908) recognized—*Arawana*, *Brumus*, and *Exochomus*—were later considered valid genera (see Gordon 1985).

Chapin (1965) proposed the new genus *Brunoides*, with the Old World *Coccinella suturalis* Fabricius designated as type, and included both *B. davisi* (Leng) and *B. septentrionis* as valid species. I am retaining the generic name *Brunoides* for *septentrionis davisi*, although Kovář (1995) restricted this genus to the Old World; he did not, however, propose a new generic name to accommodate the New World species.

Brunoides septentrionis davisi is broadly oval, convex, 3.5 to 4.5 mm long, with the upper surface glabrous. The head and pronotum are black and the elytra yellowish brown to red and coarsely punctured. Each elytron typically has two large black discal spots with the sutural area broadly black; the black areas often are confluent. Gordon (1985) illustrated the adult habitus and male and female genitalia.

DISTRIBUTION

Historical (Table 1).—State and provincial records that have been added since Leng's (1908) original description are New York (Leonard 1928), Minnesota and Wisconsin ("Lake Superior" was cited by Leng 1908) (Wingo 1952), South Carolina (Kirk 1970), and Ontario and Quebec (McNamara 1991). In his key to species of *Brunoides*, Gordon (1985) gave the general range of *B. septentrionis davisi* as "eastern United States, New York and New Jersey west to Wisconsin," but under its distribution he stated that the subspecies ranges from "southeastern Canada to Virginia, west to Minnesota." Gordon (1985) did not include Alabama, listed by Leng (1908), in the known distribution. The distribution given by Belicek (1976) for *B. septentrionis* included Illinois and Indiana, and his map appears also to include Tennessee; he did not recognize subspecies, but his records for these three states might apply to *B. septentrionis davisi*. Neither Belicek (1976) nor

Gordon (1985) listed locality data for the specimens they examined. Downie and Arnett's (1996) inclusion of Connecticut, Delaware, Maryland, Massachusetts, New Hampshire, Ohio, Pennsylvania, Rhode Island, Vermont, and Virginia in the known distribution apparently is based on their interpretation of the stippled area on Gordon's (1985) map.

New records.—*Brunoides septentrionis davisi* was collected mainly during studies on mirids (Wheeler 1991; A.G.W., unpublished data) and other hemipterans found in northeastern pitch pine-scrub oak barrens (Wheeler 1996, 1999a, b; Wheeler and Wilson 1996). Voucher specimens have been deposited in the Clemson University Arthropod Collection; Pennsylvania Department of Agriculture Collection, Harrisburg (PADA); and the National Museum of Natural History, Smithsonian Institution, Washington, D.C. (USNM).

CONNECTICUT: Hartford Co., Shaker Pines, Enfield, 28 Sept. 1991; Windham Co., Windham Airport, 1 Sept. 1991. MASSACHUSETTS: Barnstable Co., North Falmouth, 29 May 1988; Franklin Co., Montague Sand Plain, 15 June 1991. NEW HAMPSHIRE: Merrimack Co., Concord barrens, 14 Sept. 1991. NEW YORK: Suffolk Co., Yaphank, 29 May 1982; Ulster Co., Sam's Point Dwarf Pine Ridge Preserve NE of Cragmoor, 1 & 14 June and 29 Sept. 1991. NORTH CAROLINA: Rockingham Co., Rt. 220, 5.6 km S of Virginia state line, 3.3 km NNW of Stoneville, 9 Apr. 1979; Guilford Co., Rt. 68, 6.4 km S of Rt. 421, 5 km SE of Sandy Ridge, 10 April 1983. PENNSYLVANIA: Chester Co., Goat Hill Serpentine Barrens, SW of Nottingham, 8 July 1990; Lancaster Co., New Texas Serpentine Barrens, 7 July 1988; Luzerne Co., Milnesville, 17 Sept. 2001; Schuylkill Co., jct. Rt. 81 & Rt. 61, S of Frackville, 20 June & 6 Oct. 1991. SOUTH CAROLINA: Oconee Co., Rt. 76, 10.5 km NW of Westminster, 11 Nov. 2001. VIRGINIA: Botetourt Co., Eagle Rock, 8 May 1999; Rt. 81 Rest Area, 2.4 km N of

jet. Rt. 640, SE of Fincastle, 26 May & 29 July 1978, 23 May 1981, 9 June 1982, 4 Mar. & 8 Apr. 1983; Henry Co., Rt. 220, 4.4 km SW of Ridgeway, 14 Mar. 1982. WEST VIRGINIA: Mercer Co., Rt. 77, 0.6 km N of Bluestone River SE of Camp Creek, 8 May 1999.

PLANT ASSOCIATIONS AND HABITATS

Brunoides septentrionis davisi is a conifer inhabitant found almost exclusively on pines. Leng (1908) noted its occurrence on pines [likely pitch pine, *Pinus rigida* Mill.] at Lakehurst, N.J. Few specimens from the localities listed in Table 1 bear host-plant data, but those from Berlin, Framingham, and Sherborn, Mass., are labeled as found on pitch pine. At least historically, pitch pine was present at nearly all the north-eastern localities from which this coccinellid is known (e.g., Bromley 1935, Cryan 1985, Motzkin et al. 1999). The specimen from near Gordon, Wis., was taken on jack pine (*P. banksiana* Lamb.).

The adults that I collected from the New England states, New York, and Pennsylvania were from pitch pine except for a specimen from eastern red-cedar (*Juniperus virginiana* L.) in Lancaster County, Pennsylvania. Specimens from North Carolina, South Carolina, and Virginia were from Virginia pine (*P. virginiana* Mill.); the West Virginia record is based on an adult from table mountain pine (*P. pungens* Lamb.).

In the Northeast, *B. septentrionis davisi* was found in pitch pine-scrub oak barrens and similar communities. It was collected in the dwarf pine plains (ridges) near Lake Maratanza in Ulster County, New York; the Montague Sand Plain in Franklin County, Massachusetts; a pitch pine community in Schuylkill County, Pennsylvania; and remnant pine barrens such as Concord, N.H.; Shaker Pines near Entfield, Conn.; and Yaphank (Long Island), N.Y. It also was found on pitch pine in serpentine barrens in Chester and Lancaster counties, Pennsylvania. In Luzerne County, Pennsylvania, it was found on pitch pine in a ruderal site, a coal

spoilbank. Collections from table mountain pine and Virginia pine in North Carolina, South Carolina, Virginia, and West Virginia also did not involve specialized communities but were from disturbed sites, such as roadsides, in the Piedmont and in the Valley and Ridge ecoregions.

POSSIBLE PREY ASSOCIATIONS

Predation on scale insects is well known among chilocorine coccinellids (e.g., Balduf 1935, Majerus 1994, Hodek and Honěk 1996, Dixon 2000). Within *Brunoides*, the Old World *B. suturalis* preys mainly on armored scales (Diaspididae), mealybugs (Pseudococcidae), and soft scales (Coccidae) (e.g., Gautam 1990, Williams and Greathead 1990, Carnegie 1997, Ponsonby and Copland 1997), with aphids and other insects serving as alternative prey (e.g., Gautam 1990). In addition, Gordon (1985) stated that members of this genus (species unspecified) feed on cochineal or daetyllopiid scales (Daetyllopiidae) and mealybugs. For *B. septentrionis*, Belicek (1976) reported that the balsam woolly adelgid, *Adelges piceae* Ratzeburg, serves as prey, apparently referring to predation by one of the western subspecies of this lady beetle. No prey records for the easternmost subspecies, *B. septentrionis davisi*, are available, although Leng (1908) noted its occurrence at Lakehurst, N.J., on aphid-infested pines.

No definite prey records were obtained during my field work. Aphids of several genera (mainly *Cinara*) were observed at all sites where the coccinellid was found and might serve at least as alternative prey. All 10 adults of *B. septentrionis davisi* that were observed in Luzerne County, Pennsylvania, in September 2001 were on a single pitch pine growing on a coal spoilbank with 8–10 other pitch pines; the coccinellid was beaten only from branches infested with a pine needle scale, *Chionaspis heterophyllae* Cooley. In Franklin County, Massachusetts, and Schuylkill County, Pennsylvania, this lady beetle also was found on pitch pines that harbored a mar-

garrigid scale, *Matsucoccus gallicolus* Morrison. The frequent co-occurrence (ca. 5 additional sites) of the coccinellid with a more or less specialized predator of *Matsucoccus* scales (e.g., Lussier 1965), the anthecorid *Elatophilus inimicus* (Drake and Harris), suggests that *M. gallicolus* also was present at other collection sites for the coccinellid.

DISCUSSION

Brunoides septentrionis davisi is about as large as many of our familiar and well-studied coccinelline lady beetles. Moreover, its coloration—black and yellowish brown to red—is only slightly more subdued than that of our readily recognized reddish, black-spotted coccinellids. Its restriction to pines, a consistent association with specialized communities such as pitch pine-scrub oak barrens, and an apparent absence (northeastern states) on pines in Christmas tree plantations and landscape plantings might partly explain the infrequency of its collection and lack of attention given to its bionomics.

Brunoides septentrionis davisi was collected from New Hampshire to South Carolina at elevations from near sea level on Long Island, New York (17 m), to about 685 m in Ulster County, New York. Adults were found in all months from March to November (range: 4 Mar.–11 Nov.), suggesting that this coccinellid is multivoltine like most other Chilocerini (e.g., Ipert 1999).

In the Northeast, this lady beetle has been collected in several well-known pine barrens, including the New Jersey Pine Barrens and those on Cape Cod, Massachusetts, and Long Island, New York. Despite intensive collecting on pitch pines, it was not observed in New York's Albany Pine Bush or in pine barrens such as Fryeburg, Shapleigh, and Waterboro in Maine and Ossipee in New Hampshire.

In addition to the need for more field work in those northeastern pine barrens in which *B. septentrionis davisi* has yet to be found, the southern extent of its distribution

requires greater resolution. Specifically, the Alabama record, listed by Leng (1908) in describing this coccinellid but not mentioned by Gordon (1985), requires verification. It was not found in the survey of the Alabama Coccinellidae by Grimes (1965). Attempts to collect it in Alabama (and Georgia) might focus on pines in upland areas, including the Appalachian Plateau.

The taxonomic status of *B. septentrionis davisi* might also be reevaluated. Gordon (1985) indicated that he was not satisfied with his arrangement of species and subspecies of *Brunoides*. Following Leng's (1908) description of *B. septentrionis* var. *davisi*, Casey (1908) reduced this variety to a synonym of *Brunus septentrionis* Weise. According to Leng (1911), Casey, in reconsidering the status of *davisi*, thought it represented a new species that "should be called *Brunus davisi* Leng." Most subsequent authors also listed *davisi* as a valid species, including Leonard (1928) (as *Exochomus davisi*), Stehr (1930) (as *E. [Brunus] davisi*), Wingo (1952) (as *Brunus davisi*), Chapin (1965) (as *Brunoides davisi*), and Kirk (1970) (as *Brunoides davisi*). Following Gordon's (1985) monographic treatment of North American Coccinellidae, however, subsequent authors (e.g., McNamara 1991, Downie and Arnett 1996) have continued to list *davisi* as a subspecies of *B. septentrionis*.

Another gap in our knowledge of North American *Brunoides* is the lack of larval descriptions. The larval key to genera (and selected species) of Nearctic coccinellids (Rees et al. 1994) excluded this genus because larvae were unavailable for study.

The determination of this chilocerine's trophic habits also is needed. My suggestion that the diaspidid *Chionaspis heterophyllae* serves as acceptable prey is based only on the beating of 10 adults of the lady beetle from scale-infested branches of a pitch pine. The suggestion that *B. septentrionis davisi* might prey on the margarodid scale *Matsucoccus gallicolus* is offered because of the co-occurrence of these species

Table 1. Locality data for *Brunoides septentrionis davisi* from five museum collections and previously published localities (minimum of county level).

Locality	Year of Collection	Museum ^a or Reference
United States		
District of Columbia		
Washington (Rock Creek)	1901	USNM
Maryland		
Glen Echo	1992	USNM
Massachusetts		
Berkley	1936	MCZ
Berlin	1937	MCZ
Dover	1900	MCZ
Fall River	NA ^b	MCZ
Framingham	1944	MCZ
Marion	1902-1904	MCZ, USNM
Sherborn	1934	MCZ
Springfield	NA	MCZ
Tyngsboro	1893, 1908	MCZ
Michigan		
Whitefish Point	NA	USNM
Minnesota		
Duluth	1918	Stehr 1930
Hubbard County	1929	Stehr 1930
New Hampshire		
Durham	1899	UNH
New Jersey		
Browns Mill	1911	USNM
Clementon	1906	USNM
Jamesburg	NA	Smith 1910
Lakehurst	1903-1943	Leng 1908, Smith 1910; CUIIC, USNM
Manchester	NA	USNM
Milltown	NA	Smith 1910
New York		
Keeseville	1952	CNC
Lake Ronkonkoma	1922	CUIIC
Melville	1924	CUIIC
Mt. Whiteface	1922	USNM
Riverhead	1950, 1953	CUIIC
Rockaway Beach	NA	USNM
Southold	1941, 1942	CUIIC
Wyandanch	1911	Davis 1911, Leonard 1928; USNM
Yaphank	NA	Davis 1911, Leonard 1928
Pennsylvania		
Hazleton	NA	MCZ
Rhode Island		
Lonsdale	1920	MCZ
Watch Hill	1909	MCZ
South Carolina		
Long Creek	NA	Kirk 1970

Table 1. Continued.

Locality	Year of Collection	Museum ^a or Reference
Virginia		
Mt. Vernon	1911, 1960	CNC, USNM
Wisconsin		
Gordon (4 mi. east)	1952	USNM
	Canada	
Ontario		
Constance Bay	1892	CNC
Sudbury	1932, 1935	CNC

^a CNC: Canadian National Collection, Ottawa, Ont.; CUIC: Cornell University Insect Collection, Ithaca, N.Y.; MCZ: Museum of Comparative Zoology, Harvard University, Cambridge, Mass.; UNH: University of New Hampshire, Durham; USNM: National Museum of Natural History, Smithsonian Institution, Washington, D.C.

^b NA = Not Available.

and the known coccidophagous habits of other chilocorine coccinellids. Trophic relationships of a coccinellid species should not be assumed merely from its syntopy with potential prey species, and experimental studies on *B. septentrionis davisii* are needed to identify both its essential and accepted or alternative prey (*sensu* Hodek and Honěk 1996).

Leng (1908) commented on the abundance of the coccinellid at Lakehurst in the New Jersey Pine Barrens. About 75–80% of the specimens in the USNM are from this locale (R.D. Gordon, pers. comm.). I have not found this coccinellid in the New Jersey Pine Barrens during my studies of mirids (Wheeler 1991; A.G.W., unpublished data) and other insects (Wheeler 1996; 1999a, b; Wheeler and Wilson 1996). In the absence of baseline data on population trends of this lady beetle, any evidence pointing to a recent decline in its densities is tenuous. My impression, though, is that it has become increasingly difficult to find in northeastern pine barrens. Since the early 1990s, my attempts to recollect this species have been unsuccessful at several northeastern sites where 10 or more adults had been observed one or more times in 1991. Recent collecting was most intensive at Sam's Point Dwarf Pine Ridge in New York (13 Aug. 2000, 5 Sept. 2001), but also in-

cluded efforts to recollect *B. septentrionis davisii* in the pitch pine community near Frackville, Pa. (4, 6 Sept. 2001) and the Montague Sand Plain in Massachusetts (15 Aug. 1993, 6 Sept. 2001). In contrast, adults of another conifer-associated lady beetle, the coccinelline *Mulsantina picta* (Randall), were found at these and other sites during the 1990s and in 2000–2001.

The numbers of coccinellid species, especially those in crop fields, can fluctuate widely at a particular site from year to year (e.g., Foott 1974, Elliott and Kieckhefer 1990, Kieckhefer and Elliott 1990). If populations of *B. septentrionis davisii* actually have declined recently in pine barrens, hypotheses to account for its reduced densities might include competition from and intra-guild predation by Old World coccinellids that have become established in North America (e.g., Evans 1991, 2000; Elliott et al. 1996; Obrycki et al. 1998). The most abundant and widespread of these adventive species are *Coccinella septempunctata* L. and *Harmonia axyridis* (Pallas). The former coccinelline spread rapidly during the 1980s (e.g., Hoebeke and Wheeler 1980, Schaefer et al. 1987, Staines et al. 1990, Gordon and Vandenberg 1991), whereas the latter species began to assume dominance in some natural and managed systems in the 1990s (e.g., Tedders and Schaefer 1994,

Brown and Miller 1998, Colunga-Garcia and Gage 1998, Hesler et al. 2001). Although the establishment of *H. axyridis* in the eastern United States might be the result of an accidental introduction (Day et al. 1994), this predator was intentionally released numerous times between 1916 and 1985, mainly to control arboreal homopteran pests, including margarodid scales (e.g., Gordon 1985, McClure 1987). Adults and larvae of *H. axyridis* were found in the majority of the stands of pitch pine and Virginia pine that I surveyed in the late 1990s and in 2000–2001. Adults of two other adventive coccinellids—*Hippodamia variegata* (Goeze) and *Propylea quatuordecimpunctata* (L.)—were observed less often on pines (A.G.W., unpublished data).

In apple orchards of eastern West Virginia, the Old World *H. axyridis* has become the dominant coccinelline, displacing another Old World species, *C. septempunctata* (Brown and Miller 1998). It is, however, the potential adverse effects of nonindigenous coccinellids on lady beetles (and other insects) native to North America that is a cause of concern (e.g., Ehler 1990, Evans 1991, Horn 1991, Elliott et al. 1993, Wheeler and Hoebeke 1995). A detrimental effect on native coccinellids such as *Adalia bipunctata* (L.), *Coccinella novemnotata* Herbst, *C. transversoguttata richardsoni* Brown, *Cycloneda munda* (Say), and *Hippodamia convergens* Guérin-Méneville already has been suggested (Wheeler and Hoebeke 1995, Elliott et al. 1996, Wheeler and Stoops 1996, Brown and Miller 1998, Colunga-Garcia and Gage 1998). Coccinellids that have more specialized habitat requirements than those associated with agroecosystems might be especially vulnerable to competitive displacement by adventive coccinellids, their numbers perhaps declining more rapidly than those of agriculturally important lady beetles (Elliott et al. 1996). Any long-term monitoring of our native coccinellid fauna, therefore, might include not only species associated with agroecosystems but also those such as *B.*

septentrionis davisi that are found in specialized natural communities.

ACKNOWLEDGMENTS

For providing label data from specimens of the coccinellid I am grateful to Edward Becker (CNC), Richard Hoebeke (CUIC), David Furth (MCZ), Donald Chandler (UNH), and Robert Gordon and Natalia Vandenberg (USNM). Natalia Vandenberg additionally called my attention to and provided a copy of the paper by Kovář. My appreciation also is extended to Randall Schuh (American Museum of Natural History, New York) for checking for possible specimens in the AMNH, Cecil Smith (University of Georgia, Athens) for checking for material in the UGA collection, Paul Huth (Mohonk Preserve, New Paltz, N.Y.) for facilitating my access to Sam's Point Dwarf Pine Ridge Preserve, James Stimmel (PADA) for identifying the scale insects, and Peter Adler (Clemson University) for comments that improved the manuscript.

LITERATURE CITED

- Balduf, W. V. 1935. The Bionomics of Entomophagous Coleoptera. John S. Swift, St. Louis, 220 pp.
- Belicek, J. 1976. Coccinellidae of western Canada and Alaska with analyses of the transmontane zoogeographic relationships between the fauna of British Columbia and Alberta (Insecta: Coleoptera: Coccinellidae). *Quaestiones Entomologicae* 12: 283–409.
- Bromley, S. W. 1935. The original forest types of southern New England. *Ecological Monographs* 5: 61–89.
- Brown, M. W. and S. S. Miller. 1998. Coccinellidae (Coleoptera) in apple orchards of eastern West Virginia and the impact of invasion by *Harmonia axyridis*. *Entomological News* 109: 143–151.
- Carnegie, A. J. M. 1997. Sugarcane and bamboo, pp. 333–341. In Ben-Dov, Y. and C. J. Hodgson, eds. *Soft Scale Insects: Their Biology, Natural Enemies and Control*, Vol. 7B. Elsevier, Amsterdam.
- Casey, T. L. 1908. Notes on the Coccinellidae. *Canadian Entomologist* 40: 393–421.
- Chapin, E. A. 1965. The genera of the Chilocorini (Coleoptera, Coccinellidae). *Bulletin of the Museum of Comparative Zoology* 133: 227–271.
- Clausen, C. P. 1940. *Entomophagous Insects*. McGraw-Hill, London, 688 pp.
- Colunga-Garcia, M. and S. H. Gage. 1998. Arrival.

- establishment, and habitat use of the multicolored Asian lady beetle (Coleoptera: Coccinellidae) in a Michigan landscape. *Environmental Entomology* 27: 1574–1580.
- Cottrell, T. E. and K. V. Yeagan. 1998. Intraguild predation of an introduced lady beetle, *Harmonia axyridis* (Coleoptera: Coccinellidae), and a native lady beetle, *Coleomegilla maculata* (Coleoptera: Coccinellidae). *Journal of the Kansas Entomological Society* 71: 159–163.
- Cryan, J. F. 1985. Retreat in the barrens. *Defenders* 60(1): 18–29.
- Day, W. H., D. R. Prokrym, D. R. Ellis, and R. J. Chianese. 1994. The known distribution of the predator *Propylea quatuordecimpunctata* (Coleoptera: Coccinellidae) in the United States, and thoughts on the origin of this species and five other exotic lady beetles in eastern North America. *Entomological News* 105: 244–256.
- Dixon, A. F. G. 2000. *Insect Predator–Prey Dynamics: Ladybird Biology and Biological Control*. Cambridge University Press, Cambridge, 257 pp.
- Downie, N. M. and R. H. Arnett, Jr. 1996. *The Beetles of Northeastern North America, Vol. II: Polyphaga: Series Bostrichiformia through Curculionidea*. Sandhill Crane Press, Gainesville, Florida, pp. 891–1721.
- Ehler, L. E. 1990. Environmental impact of introduced biological-control agents: Implications for agricultural biotechnology, pp. 85–96. *In* Marois, J. J. and G. Bruening, eds. *Risk Assessment in Agricultural Biotechnology: Proceedings of the International Conference*. University of California Division of Agriculture and Natural Resources Publication 1928.
- Elliott, N. C. and R. W. Kieckhefer. 1990. Dynamics of aphidophagous coccinellid assemblages in small grain fields in eastern South Dakota. *Environmental Entomology* 19: 1320–1329.
- Elliott, N. C., R. W. Kieckhefer, and W. C. Kauffman. 1993. Effects of an invading coccinellid, *Coccinella septempunctata* L., on the structure of native coccinellid species assemblages in agricultural crops. *Bulletin of the Ecological Society of America* 74(2 suppl.): 226.
- Elliott, N., R. Kieckhefer, and W. Kauffman. 1996. Effects of an invading coccinellid on native coccinellids in an agricultural landscape. *Oecologia* 105: 537–544.
- Evans, E. W. 1991. Intra versus interspecific interactions of ladybeetles (Coleoptera: Coccinellidae) attacking aphids. *Oecologia* 87: 401–408.
- . 2000. Morphology of invasion: Body size patterns associated with establishment of *Coccinella septempunctata* (Coleoptera: Coccinellidae) in western North America. *European Journal of Entomology* 97: 469–474.
- Foott, W. H. 1974. Observations on Coccinellidae in corn fields in Essex County, Ontario. *Proceedings of the Entomological Society of Ontario* 104: 16–21.
- Gautam, R. D. 1990. Mass-multiplication techniques of coccinellid predator, lady bird beetle (*Brunnioides suturalis*). *Indian Journal of Agricultural Sciences* 60: 747–750.
- Gordon, R. D. 1985. The Coccinellidae (Coleoptera) of America north of Mexico. *Journal of the New York Entomological Society* 93: 1–912.
- Gordon, R. D. and N. Vandenberg. 1991. Field guide to recently introduced species of Coccinellidae (Coleoptera) in North America, with a revised key to North American genera of Coccinellini. *Proceedings of the Entomological Society of Washington* 93: 845–864.
- Grimes, W. H. 1965. *The Coccinellidae (Coleoptera) of Alabama*. Ph.D. Dissertation, Auburn University, Auburn, Alabama, 144 pp.
- Hesler, L. S., R. W. Kieckhefer, and D. A. Beck. 2001. First record of *Harmonia axyridis* (Coleoptera: Coccinellidae) in South Dakota and notes on its activity there and in Minnesota. *Entomological News* 112: 264–270.
- Hodek, I. and A. Honěk. 1996. *Ecology of Coccinellidae*. Kluwer Academic Publishers, Dordrecht, 464 pp.
- Hoebcke, E. R. and A. G. Wheeler, Jr. 1980. New distribution records of *Coccinella septempunctata* L. in the eastern United States (Coleoptera: Coccinellidae). *Coleopterists Bulletin* 34: 209–212.
- Horn, D. J. 1991. Potential impact of *Coccinella septempunctata* on endangered Lycanidae (Lepidoptera) in northwestern Ohio, USA, pp. 159–162. *In* Polgár, L., R. J. Chambers, A. F. G. Dixon, and I. Hodek, eds. *Behaviour and Impact of Aphidophaga*. SPB Academic Publishing, The Hague.
- Iperti, G. 1999. Biodiversity of predaceous Coccinellidae in relation to bioindication and economic importance. *Agriculture, Ecosystems and Environment* 74: 323–342.
- Kieckhefer, R. W. and N. C. Elliott. 1990. A 13-year survey of the aphidophagous Coccinellidae in maize fields in eastern South Dakota. *Canadian Entomologist* 122: 579–581.
- Kirk, V. M. 1970. A list of the beetles of South Carolina. Part 2—Mountain, Piedmont, and Southern Coastal Plain. *South Carolina Agricultural Experiment Station Technical Bulletin* 1038, 117 pp.
- Kovář, I. 1995. Revision of the genera *Brunnus* Muls. and *Exochomus* Redtb. (Coleoptera, Coccinellidae) of the Palaearctic Region. Part I. *Acta Entomologica Musei Nationalis Pragae* 44: 5–124.
- Leng, C. W. 1908. Notes on Coccinellidae.—III. *Journal of the New York Entomological Society* 16: 33–44.
- Leng, [C. W.]. 1911. [Proceedings of the Society, May

- 17, 1910]. Journal of the New York Entomological Society 19: 203.
- Leonard, M. D., ed. 1928. A list of the insects of New York with a list of the spiders and certain other allied groups. Cornell University Agricultural Experiment Station Memoir 101, 1121 pp.
- Lussier, S. J. 1965. A study of *Elatophilus utinica* D. & H. (Hemiptera: Anthocoridae) and its role in the natural control of the red-pine scale *Matsucoccus resinosa* B. & G. (Homoptera: Margarodidae). M.S. Thesis, University of Massachusetts, Amherst, 42 pp.
- McClure, M. S. 1987. Potential of the Asian predator, *Harmonia axyridis* Pallas (Coleoptera: Coccinellidae), to control *Matsucoccus resinosa* Bean and Godwin (Homoptera: Margarodidae) in the United States. Environmental Entomology 16: 224-230.
- McNamara, J. 1991. Family Coccinellidae, ladybird beetles, pp. 229-237. In Bousquet, Y., ed. Checklist of Beetles of Canada and Alaska. Research Branch Agriculture Canada Publication 1861/E.
- Majerus, M. E. N. 1994. Ladybirds. HarperCollins, London, 367 pp.
- Motzkin, G., W. A. Patterson III, and D. R. Foster. 1999. A historical perspective on pitch pine-scrub oak communities in the Connecticut Valley of Massachusetts. Ecosystems 2: 255-273.
- Obrycki, J. J. and T. T. Kring. 1998. Predaceous Coccinellidae in biological control. Annual Review of Entomology 43: 295-321.
- Obrycki, J. J., K. L. Giles, and A. M. Ormord. 1998. Interactions between an introduced and indigenous coccinellid species at different prey densities. Oecologia 117: 279-285.
- Ponsonby, D. J. and M. J. W. Copland. 1997. Predators, Coccinellidae and other Coleoptera, pp. 29-60. In Ben-Dov, Y. and C. J. Hodgson, eds., Soft Scale Insects: Their Biology, Natural Enemies and Control, Vol. 7B. Elsevier, Amsterdam.
- Rees, B. E., D. M. Anderson, D. Bouk, and R. D. Gordon. 1994. Larval key to genera and selected species of North American Coccinellidae (Coleoptera). Proceedings of the Entomological Society of Washington 96: 387-412.
- Schaefer, P. W., R. J. Dysart, and H. B. Specht. 1987. North American distribution of *Coccinella septempunctata* (Coleoptera: Coccinellidae) and its mass appearance in coastal Delaware. Environmental Entomology 16: 368-373.
- Staines, C. L., Jr., M. J. Rothschild, and R. B. Trumble. 1990. A survey of the Coccinellidae (Coleoptera) associated with nursery stock in Maryland. Proceedings of the Entomological Society of Washington 92: 310-313.
- Stehr, W. C. 1930. The Coccinellidae (ladybird beetles) of Minnesota (Coleoptera). University of Minnesota Agricultural Experiment Station Technical Bulletin 75, 54 pp.
- Tedders, W. L. and P. W. Schaefer. 1994. Release and establishment of *Harmonia axyridis* (Coleoptera: Coccinellidae) in the southeastern United States. Entomological News 105: 228-243.
- Wheeler, A. G., Jr. 1991. Plant bugs of *Quercus ilicifolia*: Myriads of mirids (Heteroptera) in pitch pine-scrub oak barrens. Journal of the New York Entomological Society 99: 405-440.
- . 1996. *Eremacoris depressus* Barber: Hosts, seasonality, and first New England records of a pine seed specialist (Heteroptera: Lygaeidae). Proceedings of the Entomological Society of Washington 98: 767-773.
- . 1999a. *Scaphytopius angustatus* (Osborn) (Homoptera: Cicadellidae), a leafhopper characteristic of pitch pine-scrub oak barrens. Proceedings of the Entomological Society of Washington 101: 219-220.
- . 1999b. *Phoenicocoris claricornis* and *Pino-phyltus carneolus* (Hemiptera: Miridae): Distribution and seasonality of two specialists on microsporangiate strobili of pines. Journal of the New York Entomological Society 107: 238-246.
- Wheeler, A. G., Jr. and E. R. Hoebeke. 1995. *Coccinella novemnotata* in northeastern North America: Historical occurrence and current status. Proceedings of the Entomological Society of Washington 97: 701-716.
- Wheeler, A. G., Jr. and C. A. Stoops. 1996. Status and spread of the Palearctic lady beetles *Hippodamia variegata* and *Propylea quatuordecimpunctata* (Coleoptera: Coccinellidae) in Pennsylvania, 1993-1995. Entomological News 107: 291-298.
- Wheeler, A. G., Jr. and S. W. Wilson. 1996. Planthoppers of pitch pine and scrub oak in pine barrens communities (Homoptera: Fulgoroidea). Proceedings of the Entomological Society of Washington 98: 100-108.
- Williams, J. R. and D. J. Greathead. 1990. Sugar cane, pp. 563-578. In Rosen, D., ed. Armored Scale Insects: Their Biology, Natural Enemies and Control, Vol. 4B. Elsevier, Amsterdam.
- Wingo, C. W. 1952. The Coccinellidae (Coleoptera) of the Upper Mississippi Basin. Iowa State College Journal of Science 27: 15-53.